

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization  
International Bureau



(43) International Publication Date  
5 June 2003 (05.06.2003)

PCT

(10) International Publication Number  
**WO 03/045421 A1**

- (51) International Patent Classification<sup>7</sup>: **A61K 38/17**, 45/00, G01N 33/50, A61P 3/00
- (21) International Application Number: PCT/IB02/04581
- (22) International Filing Date: 3 October 2002 (03.10.2002)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data:  
60/334,152 28 November 2001 (28.11.2001) US
- (71) Applicant (*for all designated States except US*): GENSET S.A. [FR/FR]; Intellectual Property Department, 24, rue Royale, F-75008 Paris (FR).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): DIALYNAS, Deno [US/US]; 3425 Lebon Drive, #813, San Diego, CA 92122 (US). SCALIA, Aaron [US/US]; 1691 Los Altos Rd., San Diego, CA 92109 (US). LUCAS, John [US/US]; 339 The Valley Road, Concord, MA 01742 (US). BRIGGS, Kristen [US/US]; 2454 Mango Way, Del Mar, CA 92014 (US).
- (74) Common Representative: GENSET S.A.; c/o De Luca, Giampiero, Sero International S.A., Intellectual Property, 12, Chemin des Aulx, 1228 Plan-les-Ouates (CH).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NO, NZ, OM, PH, PL, PT, RO, RU, SD, SE, SG, SI, SK, SL, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Published:  
— with international search report
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

WO 03/045421 A1

(54) Title: AGONISTS AND ANTAGONISTS OF RYZN FOR THE TREATMENT OF METABOLIC DISORDERS

(57) Abstract: The present invention relates to the field of metabolic research, in particular the discovery of compounds effective for reducing body mass and useful for treating obesity-related diseases and disorders. The obesity-related diseases or disorders envisioned to be treated by the methods of the invention include, but are not limited to, hyperlipidemia, atherosclerosis, insulin resistance, diabetes, and hypertension. In particular, the invention provides for methods of identifying and using AGONISTS and ANTAGONISTS of RYZN activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity.

## AGONISTS AND ANTAGONISTS OF RYZN FOR THE TREATMENT OF METABOLIC DISORDERS

Field of the Invention

5           The present invention relates to the field of metabolic research, in particular the discovery of compounds effective for reducing body mass and maintaining weight loss and useful for treating obesity-related diseases and disorders. The obesity-related diseases or disorders envisioned to be treated by the methods of the invention include, but are not limited to, hyperlipidemia, atherosclerosis, insulin resistance, diabetes, and hypertension. The present invention additionally  
10 relates elsewhere to the field of metabolic research, in particular the discovery of compounds effective for increasing body mass and useful for treating disorders associated with excessive weight loss. Applicant reserves the right to exclude any of the aforesaid obesity-related diseases or disorders. The disorders associated with excessive weight loss and envisioned to be treated by the methods of the invention include, but are not limited to, cachexia, cancer-related weight loss, AIDS-  
15 related weight loss, chronic inflammatory disease-related weight loss, and anorexia. Applicant reserves the right to exclude any of the aforesaid disorders associated with excessive weight loss.

In particular, the invention provides for methods of identifying and using AGONISTS and ANTAGONISTS of RYZN activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity.

20

Background of the Invention

The following discussion is intended to facilitate the understanding of the invention, but is not intended nor admitted to be prior art to the invention.

Obesity is a public health problem that is serious, widespread, and increasing. In the United States, 20 percent of the population is obese; in Europe, a slightly lower percentage is obese  
25 (Friedman (2000) Nature 404:632-634). Obesity is associated with increased risk of hypertension, cardiovascular disease, diabetes, and cancer as well as respiratory complications and osteoarthritis (Kopelman (2000) Nature 404:635-643). Even modest weight loss ameliorates these associated conditions.

Recently it was shown that particular carboxyl-terminal fragments of the full-length  
30 ACRP30 (mouse) and APM1 (human) polypeptides have unexpected effects *in vitro* and *in vivo*, including utility for weight reduction, prevention of weight gain, and control of blood glucose levels (Fruebis et al (2001) Proc Natl Acad Sci USA 98:2005-10). The effects of ACRP30 fragment administration in mammals also include reduction of elevated free fatty acid levels including elevated free fatty acid levels caused by administration of epinephrine, *i.v.* injection of "intralipid",

or administration of a high fat test meal, as well as increased fatty acid oxidation in muscle cells, and weight reduction in mammals consuming a normal or high fat/high sucrose diet.

Throughout this application, various publications, patents and published patent applications are cited. The disclosures of these publications, patents and published patent specification  
5 referenced in this application are hereby incorporated by reference into the present disclosure to more fully describe the state of the art to which this invention pertains.

#### Summary of the Invention

APM1 belongs to an expanding family of related secreted polypeptides that includes among others C2P, ZADJ-2 and ZADJ-7. These polypeptides have in common the structure: signal  
10 peptide, N-terminally disposed unique region, collagen-like region, and globular C-terminal C1q homology domain. APM1, C2P, ZADJ-2 and ZADJ-7 further share an NGLXXD amino acid motif C-terminally disposed within the globular domain within a loop implicated in receptor binding, wherein said receptor is RYZN. Fragments of APM1, C2P, ZADJ-2 and ZADJ-7 polypeptide comprising the globular domain are herein referred to as gAPM1, gC2P, gZADJ-2 and gZADJ-7. It  
15 is further taken to be understood herein that LIGAND refers to a composition consisting essentially of or consisting of *in vitro* or *in vivo* self-assembling homotrimer comprised of gAPM1, gC2P, gZADJ-2, or gZADJ-7 polypeptide fragment.

RYZN is a member of the Tumor Necrosis Factor Receptor Super Family (TNFRSF) and is a Type III transmembrane protein. The instant invention is based on RYZN as receptor for  
20 LIGAND that mediates effects, including utility for weight reduction, maintenance of weight loss, prevention of weight gain, increased insulin sensitivity, and control of blood glucose levels in humans and other mammals. These effects in mammals of RYZN engagement by LIGAND also include reduction of elevated free fatty acid levels including elevated free fatty acid levels including elevated free fatty acid levels caused by administration of epinephrine, *i.v.* injection of "intralipid",  
25 or administration of a high fat test meal, as well as increased fatty acid oxidation in muscle cells, and weight reduction in mammals consuming a normal or high fat/high sucrose diet. More specifically, the present invention is directed to RYZN to which LIGAND binds and through which LIGAND mediates said effects.

In particular, the invention provides for methods of identifying and using AGONISTS and  
30 ANTAGONISTS of RYZN activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity, as well as to pharmaceutical and physiologically acceptable compositions comprising said RYZN AGONISTS or ANTAGONISTS and methods of administering said pharmaceutical and physiologically acceptable compositions in order to increase or reduce body weight, maintain weight loss, or to treat obesity-related diseases and  
35 disorders. Assays for identifying AGONISTS and ANTAGONISTS of obesity-related activity are also part of the invention.

Preferably said RYZN AGONIST or ANTAGONIST is a compound selected from the group consisting of polypeptide, polypeptide fragment, peptide, proein, antibody, carbohydrate, lipid, small molecular weight organic compound and small molecular weight inorganic compound.

Preferably said RYZN AGONIST or ANTAGONIST is a compound that selectively binds  
5 to the extracellular domain of RYZN.

In other embodiment, said RYZN AGONIST or ANTAGONIST is a compound that selectively binds to the intracellular domain of a polypeptide comprising the extracellular domain of RYZN.

The present invention also provides a method of assaying test compounds to identify a test  
10 compound that binds to RYZN polypeptide. The method comprises contacting RYZN polypeptide with a test compound and to determine the extent of binding of the test compound to said RYZN polypeptide. The method further comprises determining whether such test compounds are AGONISTS or ANTAGONISTS of RYZN polypeptide. The present invention further provides a method of testing the impact of molecules on the expression of RYZN polypeptide or on the activity  
15 of RYZN polypeptide.

The present invention also relates to diagnostic methods of identifying individuals or non-human animals having elevated or reduced levels of RYZN products, which individuals are likely to benefit from therapies to suppress or enhance RYZN expression, respectively, and to methods of identifying individuals or non-human animals at increased risk for developing, or present state of  
20 having, certain diseases/disorders associated with RYZN abnormal expression or biological activity.

The present invention provides for methods of identifying AGONISTS of RYZN polypeptide biological activity comprising contacting a small molecule compound with RYZN polypeptides and measuring RYZN polypeptide biological activity in the presence and absence of these small molecules. The present invention further provides for methods of identifying  
25 ANTAGONISTS of RYZN polypeptide biological activity comprising contacting a small molecule compound with RYZN polypeptides and measuring RYZN polypeptide biological activity in the presence and absence of these small molecules. These small molecules can be a naturally occurring medicinal compound or derived from combinatorial chemical libraries.

The present invention also relates to pharmaceutical or physiologically acceptable  
30 compositions comprising, an active agent, including AGONIST or ANTAGONIST of the present invention.

In a first aspect, the invention is directed to RYZN AGONISTS, wherein said AGONIST is an antibody that specifically binds RYZN, a compound excluding said RYZN antibody (e.g., small organic or inorganic compound, protein, peptide, carbohydrate, lipid), or a LIGAND polypeptide or  
35 fragment thereof.

In a further preferred embodiment, the invention is directed to a RYZN AGONIST, wherein said AGONIST is an antibody that specifically binds RYZN. More preferably the invention is directed to said RYZN antibody, wherein said RYZN antibody binds RYZN and manifests LIGAND activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein.

In a further preferred embodiment, the invention is directed to a RYZN AGONIST, wherein said AGONIST is a compound excluding said RYZN antibody. More preferably the invention is directed to said compound, wherein said compound binds RYZN and manifests LIGAND activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein. Further more preferably the invention is directed to said compound, wherein said compound manifests LIGAND activity exclusive of binding to RYZN, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein. Further more preferably the invention is directed to said compound, wherein said compound increases RYZN expression.

In a further preferred embodiment, the invention is directed to a RYZN AGONIST that selectively binds to a polypeptide comprising the extracellular domain of RYZN.

In a further preferred embodiment, the invention is directed to a RYZN AGONIST, wherein said AGONIST is LIGAND, and wherein it is understood that LIGAND refers to a composition consisting essentially of or consisting of *in vitro* or *in vivo* self-assembling homotrimer comprised of gAPM1, gC2P, gZADJ-2, or gZADJ-7 polypeptide fragment. More preferably the invention is directed to said LIGAND, wherein said LIGAND binds RYZN and elicits biological activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein. More preferably the invention is directed to said LIGAND, wherein said LIGAND induces, enhances, or potentiates said biological activity exclusive of binding to RYZN. In preferred embodiment, said homotrimer is comprised of preferred gAPM1, gC2P, gZADJ-2 or gZADJ-7 polypeptide fragment.

APM1. Preferred gAPM1 polypeptide fragment is selected from amino acids 18-244, 34-244, 49-244, 56-244, 59-244, 66-244, 69-244, 78-244, 85-244, 93-244, 101-244, 102-244, 103-244, 104-244, 107-244, **110-244** or **113-244**, wherein said numbering of said amino acids within APM1 amino acid sequence is understood to be taken from said APM1 amino acid sequence presented in Table 2. Less preferred gAPM1 fragments are indicated in bold.

C2P. Preferred gC2P polypeptide fragment is selected from amino acids 20-333, 25-333, 43-333, 45-333, 46-333, 50-333, 53-333, 61-333, 67-333, 74-333, 75-333, 77-333, 81-333, 82-333, 86-333, 89-333, 95-333, 100-333, 104-333, 113-333, 116-333, 125-333, 128-333, 140-333, 160-333, 164-333, 179-333, 182-333, 185-333, 188-333, 191-333, 193-333, or **202-333**, wherein said numbering of said amino acids within C2P amino acid sequence is understood to be taken from said

C2P amino acid sequence presented in Table 2. Less preferred gC2P fragments are indicated in bold.

ZADJ-2. Preferred gZADJ-2 polypeptide fragment is selected from amino acids **16-285, 25-285, 26-285, 29-285, 30-285, 91-285, 93-285, 97-285, 98-285, 99-285, 105-285, 109-285, 112-285,**  
 5 **120-285, 126-285, 127-285, 130-285, 132-285, 133-285, 134-285, or 150-285,** wherein said numbering of said amino acids within ZADJ-2 amino acid sequence is understood to be taken from said ZADJ-2 amino acid sequence presented in Table 2. Less preferred gZADJ-2 fragments are indicated in bold.

ZADJ-7. Preferred gZADJ-7 polypeptide fragment is selected from amino acids **31-303, 39-**  
 10 **303, 78-303, 81-303, 84-303, 85-303, 88-303, 91-303, 97-303, 99-303, 109-303, 117-303, 118-303, 127-303, 139-303, 142-303, 155-303, or 162-303,** wherein said numbering of said amino acids within ZADJ-7 amino acid sequence is understood to be taken from said ZADJ-7 amino acid sequence presented in Table 2. Less preferred gZADJ-7 fragments are indicated in bold.

More preferred LIGAND is APM1.

15 In a further preferred embodiment, said AGONIST is able to lower circulating (either blood, serum or plasma) levels (concentration) of: (i) free fatty acids, (ii) glucose, and/or (iii) triglycerides.

Further preferred AGONISTS are those that significantly stimulate muscle lipid or free fatty acid oxidation as compared to untreated cells. Further preferred AGONISTS are those that cause C2C12 cells differentiated in the presence of said AGONISTS to undergo at least 10%, 20%, 30%,  
 20 35%, or 40% more oleate oxidation as compared to untreated cells.

Further preferred AGONISTS are those that increase by at least 10%, 20%, 30%, 35%, or 40% leptin uptake in a liver cell line [preferably BPRCL mouse liver cells (ATCC CRL-2217)] as compared to untreated cells.

Further preferred AGONISTS are those that significantly reduce the postprandial increase in  
 25 plasma free fatty acids or triglycerides, particularly following a high fat meal.

Further preferred AGONISTS are those that significantly reduce or eliminate ketone body production, particularly following a high fat meal.

Further preferred AGONISTS are those that increase glucose uptake in skeletal muscle cells.

Further preferred AGONISTS are those that increase glucose uptake in adipose cells.

30 Further preferred AGONISTS are those that increase glucose uptake in neuronal cells.

Further preferred AGONISTS are those that increase glucose uptake in red blood cells.

Further preferred AGONISTS are those that increase glucose uptake in the brain.

Further preferred AGONISTS are those that significantly reduce the postprandial increase in plasma glucose following a meal, particularly a high carbohydrate meal.

Further preferred AGONISTS are those that significantly prevent the postprandial increase in plasma glucose following a meal, particularly a high fat or a high carbohydrate meal.

5 Further preferred AGONISTS are those that improve insulin sensitivity.

Further preferred said AGONISTS are those that decrease body mass, wherein said decrease in body mass is comprised of a change in mass of the subcutaneous adipose tissue.

Further preferred said AGONISTS are those that decrease body mass, wherein said decrease in body mass is comprised of a change in mass of the visceral (omental) adipose tissue.

10 In a second aspect, the invention features a pharmaceutical or physiologically acceptable composition comprising, consisting essentially of, or consisting of, said AGONIST described in the first aspect and, alternatively, a pharmaceutical or physiologically acceptable diluent.

In a third aspect, the invention features a method of reducing body mass comprising providing or administering to individuals in need of reducing body mass said pharmaceutical or  
15 physiologically acceptable composition described in the second aspect.

In a fourth aspect, the invention features a method of preventing or treating an obesity-related disease or disorder comprising providing or administering to an individual in need of such treatment said pharmaceutical or physiologically acceptable composition described in the second aspect. Preferably, said obesity-related disease or disorder is selected from the group consisting of  
20 obesity, insulin resistance, atherosclerosis, atheromatous disease, heart disease, hypertension, stroke, Syndrome X, Noninsulin Dependent Diabetes Mellitus (NIDDM, or Type II diabetes) and Insulin Dependent Diabetes Mellitus (IDDM or Type I diabetes). Diabetes-related complications to be treated by the methods of the invention include microangiopathic lesions, ocular lesions, retinopathy, neuropathy, and renal lesions. Heart disease includes, but is not limited to, cardiac  
25 insufficiency, coronary insufficiency, and high blood pressure. Other obesity-related disorders to be treated by said RYZN AGONIST of the invention include hyperlipidemia and hyperuricemia. In preferred embodiments, said individual is a mammal, preferably a human.

In related aspects, embodiments of the present invention includes methods of causing or inducing a desired biological response in an individual comprising the steps of: providing or  
30 administering to an individual a composition comprising AGONIST, wherein said biological response is selected from the group consisting of:

- (a) lowering circulating (either blood, serum, or plasma) levels (concentration) of free fatty acids;

- (b) lowering circulating (either blood, serum or plasma) levels (concentration) of glucose;
- (c) lowering circulating (either blood, serum or plasma) levels (concentration) of triglycerides;
- 5 (d) stimulating muscle lipid or free fatty acid oxidation;
- (e) increasing leptin uptake in the liver or liver cells;
- (e) reducing the postprandial increase in plasma free fatty acids, particularly following a high fat meal;
- (f) reducing or eliminating ketone body production, particularly following a high fat meal;
- 10 (g) increasing tissue sensitivity to insulin, particularly muscle, adipose, liver or brain,

and further wherein said biological response is significantly greater than, or at least 10%, 20%, 30%, 35%, or 40% greater than that observed in the absence of treatment; or alternatively wherein said biological response is greater than a transient response; or alternatively wherein said biological response is sustained. In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to control blood glucose in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) in combination with insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to control blood glucose in some persons with Insulin Dependent Diabetes Mellitus (IDDM, Type I diabetes) in combination with insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to control body weight in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) in combination with insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to control body weight in some persons with Insulin Dependent Diabetes Mellitus (IDDM, Type I diabetes) in combination with insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to control blood glucose in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) alone, without combination of insulin therapy.



In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to control blood glucose in some persons with Insulin Dependent Diabetes Mellitus (IDDM, Type I diabetes) alone, without combination of insulin therapy.

- 5 In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to control body weight in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) alone, without combination of insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or  
10 physiologically acceptable composition can be used as a method to control body weight in some persons with Insulin Dependent Diabetes Mellitus (IDDM, Type I diabetes) alone, without combination of insulin therapy.

In a further preferred embodiment, the present invention may be used in complementary therapy of NIDDM patients to improve their weight or glucose control in combination with an  
15 insulin secretagogue or an insulin sensitising agent. Preferably, the insulin secretagogue is 1,1-dimethyl-2-(2-morpholino phenyl)guanidine fumarate (BTS67582) or a sulphonylurea selected from tolbutamide, tolazamide, chlorpropamide, glibenclamide, glimepiride, glipizide and glidazide. Preferably, the insulin sensitising agent is selected from metformin, ciglitazone, troglitazone and pioglitazone.

- 20 The present invention further provides a method of improving the body weight or glucose control of NIDDM patients alone, without an insulin secretagogue or an insulin sensitising agent.

In a further preferred embodiment, the present invention may be used in complementary therapy of IDDM patients to improve their weight or glucose control in combination with an insulin secretagogue or an insulin sensitising agent. Preferably, the insulin secretagogue is 1,1-dimethyl-2-  
25 (2-morpholino phenyl) guanidine fumarate (BTS67582) or a sulphonylurea selected from tolbutamide, tolazamide, chlorpropamide, glibenclamide, glimepiride, glipizide and glidazide. Preferably, the insulin sensitising agent is selected from metformin, ciglitazone, troglitazone and pioglitazone.

- The present invention further provides a method of improving the body weight or glucose  
30 control of IDDM patients alone, without an insulin secretagogue or an insulin sensitising agent.

In a further preferred embodiment, the present invention may be administered either concomitantly or concurrently, with the insulin secretagogue or insulin sensitising agent for example in the form of separate dosage units to be used simultaneously, separately or sequentially (either before or after the secretagogue or either before or after the sensitising agent). Accordingly, the  
35 present invention further provides for a composition of pharmaceutical or physiologically acceptable

composition and an insulin secretagogue or insulin sensitising agent as a combined preparation for simultaneous, separate or sequential use for the improvement of body weight or glucose control in NIDDM or IDDM patients.

In further preferred embodiments, the present invention of said pharmaceutical or  
5 physiologically acceptable composition further provides a method for the use as an insulin sensitiser.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to improve insulin sensitivity in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) in combination with insulin therapy.

10 In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method to improve insulin sensitivity in some persons with Insulin Dependent Diabetes Mellitus (IDDM, Type I diabetes) in combination with insulin therapy.

In further preferred embodiments, the present invention of said pharmaceutical or  
15 physiologically acceptable composition can be used as a method to improve insulin sensitivity in some persons with Noninsulin Dependent Diabetes Mellitus (NIDDM, Type II diabetes) without insulin therapy.

In a fifth aspect, the invention features a use of AGONIST described in the first aspect for treatment of obesity-related diseases and disorders and/or reducing body mass. Preferably, said  
20 obesity-related diseases and disorders are selected from the group consisting of obesity, insulin resistance, atherosclerosis, atheromatous disease, heart disease, hypertension, stroke, Syndrome X, Noninsulin Dependent Diabetes Mellitus (NIDDM, or Type II diabetes) and Insulin Dependent Diabetes Mellitus (IDDM or Type I diabetes). Diabetes-related complications to be treated by the methods of the invention include microangiopathic lesions, ocular lesions, retinopathy, neuropathy,  
25 and renal lesions. Heart disease includes, but is not limited to, cardiac insufficiency, coronary insufficiency, and high blood pressure. Other obesity-related disorders to be treated by said AGONIST of the invention include hyperlipidemia and hyperuricemia.

In a sixth aspect, the invention features a use of AGONIST described in the first aspect for the preparation of a medicament for the treatment of obesity-related diseases and disorders and/or  
30 for reducing body mass. Preferably, said obesity-related disease or disorder is selected from the group consisting of obesity, insulin resistance, atherosclerosis, atheromatous disease, heart disease, hypertension, stroke, Syndrome X, Noninsulin Dependent Diabetes Mellitus (NIDDM, or Type II diabetes) and Insulin Dependent Diabetes Mellitus (IDDM or Type I diabetes). Diabetes-related complications to be treated by the methods of the invention include microangiopathic lesions, ocular  
35 lesions, retinopathy, neuropathy, and renal lesions. Heart disease includes, but is not limited to, cardiac insufficiency, coronary insufficiency, and high blood pressure. Other obesity-related

disorders to be treated by compounds of the invention include hyperlipidemia and hyperuricemia. In preferred embodiments, said individual is a mammal, preferably a human.

In a seventh aspect, the invention provides AGONIST of the first aspect of the invention, or a composition of the second aspect of the invention, for use in a method of treatment of the human  
5 or animal body.

In an eighth aspect, the invention features methods of reducing body weight comprising providing to an individual said pharmaceutical or physiologically acceptable composition described in the second aspect, or AGONIST described in the first aspect. Where the reduction of body weight is practiced for cosmetic purposes, the individual has a BMI of at least 20 and no more than 25. In  
10 embodiments for the treatment of obesity, the individual may have a BMI of at least 20. One embodiment for the treatment of obesity provides for the treatment of individuals with BMI values of at least 25. Another embodiment for the treatment of obesity provides for the treatment of individuals with BMI values of at least 30. Yet another embodiment provides for the treatment of individuals with BMI values of at least 40.

15 In further embodiment, the invention features methods of maintaining weight loss comprising providing to an individual said pharmaceutical or physiologically acceptable composition.

In a ninth aspect, the invention features the pharmaceutical or physiologically acceptable composition described in the second aspect for reducing body mass and/or for treatment or  
20 prevention of obesity-related diseases or disorders. Preferably, said obesity-related disease or disorder is selected from the group consisting of obesity, insulin resistance, atherosclerosis, atheromatous disease, heart disease, hypertension, stroke, Syndrome X, Noninsulin Dependent Diabetes Mellitus (NIDDM, or Type II diabetes) and Insulin Dependent Diabetes Mellitus (IDDM or Type I diabetes). Diabetes-related complications to be treated by the methods of the invention  
25 include microangiopathic lesions, ocular lesions, retinopathy, neuropathy, and renal lesions. Heart disease includes, but is not limited to, cardiac insufficiency, coronary insufficiency, and high blood pressure. Other obesity-related disorders to be treated by compounds of the invention include hyperlipidemia and hyperuricemia. In preferred embodiments, said individual is a mammal, preferably a human. In preferred embodiments, the identification of said individuals to be treated  
30 with said pharmaceutical or physiologically acceptable composition comprises genotyping LIGAND single nucleotide polymorphisms (SNPs) or measuring LIGAND polypeptide or mRNA levels in clinical samples from said individuals. Preferably, said clinical samples are selected from the group consisting of blood, serum, plasma, urine, and saliva.

In a tenth aspect, the invention features the pharmaceutical or physiologically acceptable  
35 composition described in the second aspect for reducing body weight for cosmetic reasons.

In an eleventh aspect, AGONIST of the invention is used in methods of treating insulin resistance comprising providing to an individual said pharmaceutical or physiologically acceptable composition described in the second aspect, or AGONIST described in the first aspect.

In a preferred aspect of the methods above and disclosed herein, the amount of AGONIST  
5 administered to an individual is sufficient to bring levels of RYZN activation to their normal levels (levels in individuals without obesity-related disease or disorder). "Normal levels" of RYZN activation may be followed using surrogate markers including circulating (either blood, serum or plasma) levels (concentration) of: (i) free fatty acids, (ii) glucose, and/or (iii) triglycerides.

In a twelfth aspect, the invention is directed to a RYZN ANTAGONIST, wherein said  
10 ANTAGONIST is a soluble fragment of RYZN polypeptide, an antibody that specifically binds RYZN, a compound excluding said soluble fragment of RYZN polypeptide and said RYZN antibody (e.g., small molecular weight organic or inorganic compound, protein, peptide, carbohydrate, lipid), or a variant or fragment of LIGAND polypeptide.

In a further preferred embodiment, the invention is directed to a RYZN ANTAGONIST,  
15 wherein said ANTAGONIST is a soluble fragment of RYZN polypeptide. More preferably the invention is directed to purified, isolated, or recombinant soluble fragments of RYZN polypeptide. More preferably the invention is directed to said soluble fragment of RYZN polypeptide, wherein said soluble fragment binds LIGAND and blocks LIGAND activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or  
20 described herein, and wherein said soluble fragment of RYZN polypeptide does not activate RYZN. Preferably said soluble fragment of RYZN polypeptide blocks or inhibits LIGAND binding to RYZN. In preferred embodiments, said soluble fragment of RYZN polypeptide comprises, consists essentially of, or consists of, at least 6 and not more than 113 consecutive amino acids of SEQ ID NO:2, more preferably of amino acids comprising the extracellular domain of RYZN. Preferred said  
25 soluble fragment of RYZN comprises the extracellular domain of mature RYZN polypeptide. Particularly preferred soluble fragment of RYZN comprises amino acids 1-61, 2-61, 1-63, 2-63, 1-68, 2-68, 1-73, 2-73, 1-74, 2-74, 1-76, 2-76, 1-89, 2-89, 1-96, 2-96, 1-108, 2-108, 1-113 or 2-113 of SEQ ID NO:2. In other preferred embodiments, said soluble fragment of RYZN polypeptide comprises an amino acid sequence at least 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%,  
30 98%, or 99% identical to the corresponding consecutive amino acids of SEQ ID NO:2. Further preferred embodiments include heterologous polypeptides comprising a RYZN polypeptide of the invention. In further preferred embodiment, a RYZN polypeptide of the invention is conjugated at its N- or C-terminus to an antibody Fc region or portion thereof.

In a further preferred embodiment, the invention is directed to a RYZN ANTAGONIST,  
35 wherein said ANTAGONIST is an antibody that specifically binds RYZN. More preferably the invention is directed to said RYZN antibody, wherein said RYZN antibody binds RYZN and blocks

LIGAND activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein, and wherein said RYZN antibody does not activate RYZN. Preferably said RYZN antibody blocks or inhibits LIGAND binding to RYZN.

In a further preferred embodiment, the invention is directed to a RYZN ANTAGONIST,  
5 wherein said ANTAGONIST is a compound excluding said soluble fragment of RYZN polypeptide and said RYZN antibody (e.g., small organic molecule, protein, peptide). More preferably the invention is directed to said compound, wherein said compound binds to RYZN and blocks LIGAND activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein, and wherein said compound does not  
10 activate RYZN. Preferably said compound that binds to RYZN blocks or inhibits LIGAND binding to RYZN. Further more preferably the invention is directed to said compound, wherein said compound blocks or inhibits LIGAND activity exclusive of binding to RYZN, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein, and wherein said compound does not activate RYZN. Further more preferably the  
15 invention is directed to said compound, wherein said compound blocks or inhibits RYZN expression and wherein said compound does not have LIGAND activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein, and wherein said compound does not activate RYZN.

In a further preferred embodiment, the invention is directed to a RYZN ANTAGONIST,  
20 wherein said ANTAGONIST is a variant or fragment of LIGAND polypeptide. More preferably the invention is directed to said variant or fragment of LIGAND polypeptide, wherein said variant or fragment of LIGAND polypeptide binds RYZN and blocks LIGAND activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or described herein, and wherein said variant or fragment of LIGAND polypeptide does not activate  
25 RYZN. Preferably said variant or fragment of LIGAND polypeptide blocks or inhibits LIGAND binding to RYZN. More preferably the invention is directed to said variant or fragment of LIGAND polypeptide, wherein said variant or fragment of LIGAND polypeptide inhibits the induction, enhancement, or potentiation of said biological activity exclusive of binding to RYZN.

In a further preferred embodiment, the invention is directed to a RYZN ANTAGONIST that  
30 selectively binds to a polypeptide comprising the extracellular domain of RYZN.

In a further preferred embodiment, said ANTAGONIST is able to raise circulating (either blood, serum or plasma) levels (concentration) of: (i) free fatty acids, (ii) glucose, and/or (iii) triglycerides.

Further preferred said ANTAGONISTS are those that significantly inhibit muscle lipid or  
35 free fatty acid oxidation stimulated by its LIGAND. Further preferred said ANTAGONISTS are

those that cause C2C12 cells differentiated in the presence of LIGAND to undergo at least 10%, 20%, 30%, 35%, or 40% less oleate oxidation as compared to untreated cells.

Further preferred said ANTAGONISTS are those that inhibit by at least 10%, 20%, 30%, 35%, or 40% the increase in leptin uptake stimulated by LIGAND polypeptide in a liver cell line  
5 [preferably BPRCL mouse liver cells (ATCC CRL-2217)] as compared to untreated cells.

Further preferred said ANTAGONISTS are those that significantly increase the postprandial increase in plasma free fatty acids, particularly following a high fat meal.

Further preferred said ANTAGONISTS are those that significantly increase ketone body production, particularly following a high fat meal.

10 Further preferred said ANTAGONISTS are those that decrease glucose uptake in skeletal muscle cells stimulated by LIGAND.

Further preferred said ANTAGONISTS are those that decrease glucose uptake in adipose cells stimulated by LIGAND.

Further preferred said ANTAGONISTS are those that decrease glucose uptake in neuronal  
15 cells stimulated by LIGAND.

Further preferred said ANTAGONISTS are those that decrease glucose uptake in red blood cells stimulated by LIGAND.

Further preferred said ANTAGONISTS are those that decrease glucose uptake in the brain stimulated by LIGAND.

20 Further preferred said ANTAGONISTS are those that significantly increase the postprandial increase in plasma glucose following a meal, particularly a high carbohydrate meal.

Further preferred said ANTAGONISTS are those that significantly facilitate the postprandial increase in plasma glucose following a meal, particularly a high fat or a high carbohydrate meal.

25 Further preferred said ANTAGONISTS are those that reduce the insulin sensitivity stimulated by LIGAND.

Further preferred said ANTAGONISTS are those that increase body mass, wherein said increase in body mass is comprised of a change in mass of the subcutaneous adipose tissue.

Further preferred said ANTAGONISTS are those that increase body mass, wherein said  
30 increase in body mass is comprised of a change in mass of the visceral (omental) adipose tissue.

In a thirteenth aspect, the invention features a pharmaceutical or physiologically acceptable composition comprising, consisting essentially of, or consisting of, said ANTAGONIST described in the twelfth aspect and, alternatively, a pharmaceutical or physiologically acceptable diluent.

In a fourteenth aspect, the invention features a method of increasing body mass comprising providing or administering to individuals in need of increasing body mass said pharmaceutical or physiologically acceptable composition described in the thirteenth aspect.

In a fifteenth aspect, the invention features a method of preventing or treating disorders  
5 associated with excessive weight loss comprising providing or administering to an individual in need of such treatment said pharmaceutical or physiologically acceptable composition described in the thirteenth aspect. Preferably said disorder is selected from the group consisting of cachexia, wasting, cancer-related weight loss, AIDS-related weight loss, chronic inflammatory disease-related weight loss, anorexia, and bulimia. Said disorders associated with excessive weight loss are  
10 comprised of those mediated by tumor necrosis factor (TNFalpha) alone, those mediated by TNFalpha plus one or more additional factors, and those mediated only by one or more factors exclusive of TNFalpha. Said factors include, but are not restricted to, macrophage migration inhibitory factor, interleukin 1, and interleukin 6. In preferred embodiments, said individual is a mammal, preferably a human.

15 In related aspects, embodiments of the present invention includes methods of causing or inducing a desired biological response in an individual comprising the steps of: providing or administering to an individual a composition comprising ANTAGONIST, wherein said biological response is selected from the group consisting of:

- 20 (a) raising circulating (either blood, serum, or plasma) levels (concentration) of free fatty acids (FFA) or triglycerides (TG);
- (b) raising circulating (either blood, serum or plasma) levels (concentration) of glucose;
- (c) raising circulating (either blood, serum or plasma) levels (concentration) of triglycerides;
- (d) inhibiting muscle lipid or free fatty acid oxidation;
- 25 (c) inhibiting leptin uptake in the liver or liver cells;
- (e) increasing the postprandial increase in plasma free fatty acids, particularly following a high fat meal; and,
- (f) increasing or eliminating ketone body production, particularly following a high fat meal;
- 30 (g) reducing tissue sensitivity to insulin, particularly muscle, adipose, liver or brain,

and further wherein said biological response is greater than a transient response; or alternatively wherein said biological response is sustained. In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition can be used as a method

of increasing body mass in some persons with cachexia, wasting, cancer-related weight loss, AIDS-related weight loss, chronic inflammatory disease-related weight loss, anorexia, and bulimia.

In further preferred embodiments, the present invention of said pharmaceutical or physiologically acceptable composition further provides a method for the use as an insulin de-  
5 sensitiser, wherein the sensitivity of a cell or tissue to insulin is reduced.

In a sixteenth aspect, the invention features a method of making the RYZN polypeptide described in the twelfth aspect, wherein said method is selected from the group consisting of proteolytic cleavage, recombinant methodology and artificial synthesis. In a preferred embodiment, proteolytic cleavage is carried out using trypsin, plasmin, or collagenase.

10 In a seventeenth aspect, the invention features a use of ANTAGONIST described in the twelfth aspect for the preparation of a medicament for the treatment of disorders associated with excessive weight loss and/or for increasing body mass. Preferably, said disorder is selected from the group consisting of cachexia, wasting, cancer-related weight loss, AIDS-related weight loss, chronic inflammatory disease-related weight loss, anorexia, and bulimia. In preferred embodiments, said  
15 individual is a mammal, preferably a human.

In an eighteenth aspect, the invention provides ANTAGONIST of the twelfth aspect of the invention, or a composition of the thirteenth aspect of the invention, for use in a method of treatment of the human or animal body.

In a nineteenth aspect, the invention features methods of increasing body weight comprising  
20 providing to an individual said pharmaceutical or physiologically acceptable composition described in the thirteenth aspect, or ANTAGONIST described in the twelfth aspect. Where the increase of body weight is practiced for cosmetic purposes, the individual has a BMI of no greater than 25 and at least 20. In embodiments for the treatment of disorders associated with excessive weight loss, the individual may have a BMI no greater than 20. One embodiment for the treatment of disorders  
25 associated with excessive weight loss provides for the treatment of individuals with BMI values of no greater than 15. Alternatively, for increasing the body weight of an individual, the BMI value should be at least 15 and no more than 20.

In a twentieth aspect, the invention features the pharmaceutical or physiologically acceptable composition described in the thirteenth aspect for increasing body mass and/or for  
30 treatment of disorders associated with excessive weight loss. Preferably, said disorder is selected from the group consisting of cachexia, wasting, cancer-related weight loss, AIDS-related weight loss, chronic inflammatory disease-related weight loss, anorexia, and bulimia. In preferred embodiments, said individual is a mammal, preferably a human.



In a twenty-first aspect, the invention features the pharmaceutical or physiologically acceptable composition described in the thirteenth aspect for increasing body weight for cosmetic reasons.

In a preferred aspect of the methods above and disclosed herein, the amount of  
5 ANTAGONIST administered to an individual is sufficient to bring levels of RYZN activation to their normal levels (levels in healthy individuals). "Normal levels" of RYZN activation may be followed using surrogate markers including circulating (either blood, serum or plasma) levels (concentration) of: (i) free fatty acids, (ii) glucose, and/or (iii) triglycerides.

#### Brief Description of Tables

10 Table 1 lists known or predicted biologic structural and functional domains for the RYZN polypeptide of SEQ ID NO:2 of the present invention, including the extracellular (EC) domain, transmembrane domain, and intracellular (IC) domain.

Table 2 lists the amino acid sequence of full-length APM1, C2P, ZADJ-2 and ZADJ-7 polypeptide. The total number of amino acids is given in parentheses. The predicted signal peptide  
15 is indicated in bold. The collagen-like region is indicated by dotted line. The region between the predicted signal peptide and the collagen-like region is the N-terminally disposed unique region. The globular C-terminal C1q homology domain is indicated by single underline. The NGLXXD amino acid motif C-terminally disposed within the globular domain is indicated by double underline. It is taken to be understood that C2P herein encompasses variants comprising the substitution of  
20 valine for methionine at position 219 and/or the substitution of methionine for valine at position 301.

#### Structure of RYZN Polypeptide

The full-length RYZN polypeptide is comprised of at least 3 distinct regions including:

1. an extracellular domain comprising a LIGAND binding portion and comprising amino acids from about amino acids 1-113 of SEQ ID NO:2;
- 25 2. a transmembrane domain comprising amino acids from about amino acids 114-136 of SEQ ID NO:2; and
3. an intracellular domain comprising amino acids from about amino acids 137-247 of SEQ ID NO:2.

#### Brief Description of Sequence Listing

30 SEQ ID NO:1 is the nucleotide sequence of cDNA with an open reading frame which location is indicated as features. When appropriate, the locations of the potential polyadenylation site and polyadenylation signal are also indicated.

SEQ ID NO:2 is the amino acid sequence of polypeptide encoded by the cDNA of SEQ ID NO:1.

The appended Sequence Listing is hereby incorporated by reference in its entirety.

### Detailed Description

#### Definitions

Before describing the invention in greater detail, the following definitions are set forth to  
5 illustrate and define the meaning and scope of the terms used to describe the invention herein.

The term "isolated" requires that the material be removed from its original environment (e.  
g., the natural environment if the material is naturally occurring).

The term "purified" does not require absolute purity; rather, it is intended as a relative  
definition. Purification of starting material or natural material to at least one order of magnitude,  
10 preferably two or three orders, and more preferably four or five orders of magnitude is expressly  
contemplated.

As used interchangeably herein, the term "polynucleotide(s)" include RNA or DNA (either  
single or double stranded, coding, complementary or antisense), or RNA/DNA hybrid sequences of  
more than one nucleotide in either single chain or duplex form (although each of the above species  
15 may be particularly specified).

The terms "complementary" or "complement thereof" are used herein to refer to the  
sequences of polynucleotides that are capable of forming Watson & Crick base pairing with another  
specified polynucleotide throughout the entirety of the complementary region.

The terms "polypeptide" and "protein", used interchangeably herein, refer to a polymer of  
20 amino acids without regard to the length of the polymer; thus, peptides, oligopeptides, and proteins  
are included within the definition of polypeptide. This term also does not specify or exclude  
chemical or post-expression modifications of the polypeptides of the invention, although chemical or  
post-expression modifications of these polypeptides may be included excluded as specific  
embodiments.

As used herein, the terms "recombinant polynucleotide" and "polynucleotide construct" are  
used interchangeably to refer to linear or circular, purified or isolated polynucleotides that have been  
artificially designed and which comprise at least two nucleotide sequences that are not found as  
contiguous nucleotide sequences in their initial natural environment. In particular, these terms mean  
that the polynucleotide or cDNA is adjacent to "backbone" nucleic acid to which it is not adjacent in  
30 its natural environment.

The term "recombinant polypeptide" is used herein to refer to polypeptides that have been  
artificially designed and which comprise at least two polypeptide sequences that are not found as  
contiguous polypeptide sequences in their initial natural environment, or to refer to polypeptides  
which have been expressed from a recombinant polynucleotide.

As used herein, the term “operably linked” refers to a linkage of polynucleotide elements in a functional relationship.

As used herein, the term “non-human animal” refers to any non-human animal, including insects, birds, rodents and more usually mammals. Both the terms “animal” and “mammal”  
5 expressly embrace human subjects unless preceded with the term “non-human”.

The term “domain” refers to an amino acid fragment with specific biological properties. This term encompasses all known structural and linear biological motifs.

As used herein, the term “receptor” refers to a polypeptide to which a “ligand” binds and through which said “ligand” elicits a biological response comprised of biological activities. Said  
10 receptor is preferably RYZN of the present invention. Said “ligand” is preferably LIGAND of the present invention. By “receptor activation” is intended “ligand”-mediated alteration of said receptor polypeptide, wherein said alteration is selected from but not limited to the group consisting of receptor alterations associated with said biological response.

As used herein, the term “AGONIST” refers to naturally occurring and synthetic compounds  
15 capable of inducing, enhancing, or potentiating a biological response comprised of biological activities.

As used herein, the term “ANTAGONIST” refers to naturally occurring and synthetic compounds capable of inhibiting a biological response, inhibiting the induction of a biological response, or inhibiting the potentiation of a biological response, wherein said biological response is  
20 comprised of biological activities.

Without being limited by theory, the compounds/polypeptides of the invention are capable of modulating the partitioning of dietary lipids between the liver and peripheral tissues, and are thus believed to treat “diseases involving the partitioning of dietary lipids between the liver and peripheral tissues.” The term “peripheral tissues” is meant to include muscle and adipose tissue. In  
25 preferred embodiments, the compounds/polypeptides of the invention partition the dietary lipids toward or away from the muscle. In alternative preferred embodiments, the dietary lipids are partitioned toward or away from the adipose tissue. In other preferred embodiments, the dietary lipids are partitioned toward or away from the liver. In yet other preferred embodiments, the compounds/polypeptides of the invention increase or decrease the oxidation of dietary lipids,  
30 preferably free fatty acids (FFA) by the muscle. Dietary lipids include, but are not limited to triglycerides and free fatty acids.

Preferred diseases believed to involve the partitioning of dietary lipids include obesity-related diseases and disorders such as obesity, insulin resistance, atherosclerosis, atheromatous disease, heart disease, hypertension, stroke, Syndrome X, Noninsulin Dependent Diabetes Mellitus  
35 (NIDDM, or Type II diabetes) and Insulin Dependent Diabetes Mellitus (IDDM or Type I diabetes).

Diabetes-related complications to be treated by the methods of the invention include microangiopathic lesions, ocular lesions, retinopathy, neuropathy, and renal lesions. Heart disease includes, but is not limited to, cardiac insufficiency, coronary insufficiency, and high blood pressure. Other obesity-related disorders to be treated by compounds of the invention include hyperlipidemia  
5 and hyperuricemia. Yet other disorders of the invention include disorders associated with excessive weight loss such as cachexia, wasting, cancer-related weight loss, AIDS-related weight loss, chronic inflammatory disease-related weight loss, anorexia, and bulimia.

The terms "comprising", "consisting of" and "consisting essentially of" may be interchanged for one another throughout the instant application, although each retains its normal definition. The  
10 term "having" has the same meaning as "comprising" and may be replaced with either the term "consisting of" or "consisting essentially of".

#### Polypeptides of the Invention

Preferably, polypeptides of the invention are recombinantly produced using routine expression methods known in the art. The polynucleotide encoding the desired polypeptide is  
15 operably linked to a promoter into an expression vector suitable for any convenient host. Both eukaryotic and prokaryotic host systems are used in forming recombinant polypeptides. The polypeptide is then isolated from lysed cells or from the culture medium and purified to the extent needed for its intended use.

Consequently, a further embodiment of the present invention is a method of making a  
20 polypeptide, said method comprising the steps of

- a) obtaining a cDNA encoding said polypeptide;
- b) inserting said cDNA in an expression vector such that the cDNA is operably linked to a promoter; and
- c) introducing said expression vector into a host cell whereby said host cell produces  
25 said polypeptide.

In one aspect of this embodiment, the method further comprises the step of isolating the polypeptide. Another embodiment of the present invention is a polypeptide obtainable by the method described in the preceding paragraph.

The expression vector is any of the mammalian, yeast, insect or bacterial expression systems  
30 known in the art. Commercially available vectors and expression systems are available from a variety of suppliers including Genetics Institute (Cambridge, MA), Stratagene (La Jolla, California), Promega (Madison, Wisconsin), and Invitrogen (San Diego, California). In preferred embodiment, recombinant polypeptides of the invention are expressed in mammalian cells.

The invention is drawn, *inter alia*, to isolated, purified or recombinant polypeptides. RYZN polypeptides of the invention are useful for increasing (ANTAGONISTS of RYZN) body weight either as a cosmetic treatment or for treatment or prevention of diseases and disorders as discussed or described herein. RYZN polypeptides are also useful *inter alia* in screening assays for  
5 AGONISTS or ANTAGONISTS of RYZN activity and for raising RYZN-specific antibodies. When used for cosmetic treatments, or for the treatment or prevention of diseases, disorders, or conditions, one or more RYZN polypeptides can be provided to a subject. Thus, various fragments of the full-length protein can be combined into a "cocktail" for use in the various treatment regimens. LIGAND polypeptides of the invention are useful for reducing (AGONISTS of RYZN)  
10 body weight either as a cosmetic treatment or prevention of diseases and disorders as discussed or described herein.

The RYZN polypeptides of the present invention are preferably provided in an isolated form, and may be partially or substantially purified.

#### Modifying RYZN biological activity

15 Modifying endogenous RYZN biological activity is expressly contemplated by the present invention. The present invention further relates to compounds able to modulate RYZN biological activity and methods to use these compounds. Such compounds may interact with RYZN polypeptides directly or indirectly.

#### Candidate AGONISTS and ANTAGONISTS Obtained by Optical Biosensor Methods

20 Compounds interacting with a polypeptide comprising RYZN extracellular domain can be screened by using an Optical Biosensor as described in Edwards and Leatherbarrow (1997) and also in Szabo *et al.* (1995), the disclosures of which are incorporated by reference. This technique permits the detection of interactions between molecules in real time, without the need of labeled molecules. This technique, which is based on the surface plasmon resonance (SPR) phenomenon, is  
25 presented in more detail in Example 1.

#### Compounds Modulating RYZN Biological Activity

Another method of screening for compounds that modulate RYZN biological activity is by measuring the effects of test compounds on specific biological activity, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or  
30 as described herein, in a host cell. In one embodiment, the present invention relates to a method of identifying an agent that alters RYZN activity, wherein a nucleic acid construct comprising the polynucleotide of SEQ ID NO:1 or a fragment thereof encoding full-length RYZN polypeptide is introduced into a mammalian host cell. The transfected mammalian host cells are maintained under conditions appropriate for expression of the encoded RYZN, whereby the nucleic acid is expressed.  
35 The host cells are then contacted with a compound to be assessed (an agent) and an activity of the

cells is detected in the presence of the compound to be assessed, wherein said activity is selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or as described herein. Detection of a change in said activity for said transfected host cell, but not in untransfected host cell, in the presence of the agent indicates that the agent alters RYZN activity. In  
5 a particular embodiment, the invention relates to a method of identifying an agent which is an activator (AGONIST) of RYZN activity, wherein detection of an increase of said activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or as described herein, in the presence of the agent indicates that the agent activates RYZN activity. In another particular embodiment, the invention relates to a method of identifying  
10 an agent which is an inhibitor (ANTAGONIST) of RYZN activity, wherein detection of a decrease of said activity, said activity being selected from the group consisting of lipid partitioning, lipid metabolism, and insulin-like activity or as described herein, in the presence of the agent indicates that the agent inhibits RYZN activity.

Detection of a change in said RYZN activity, said activity being selected from the group  
15 consisting of lipid partitioning, lipid metabolism, and insulin-like activity or as described herein, can be performed using a variety of techniques as described for representative activities in Examples provided herein.

In a particular embodiment a high throughput screen can be used to identify agents that activate (enhance) or inhibit RYZN activity (See e.g., PCT publication WO 98/45438, which  
20 disclosure is hereby incorporated by reference in its entirety).

#### Methods of Screening for Compounds Modulating RYZN Activity

The present invention also relates to methods of screening compounds for their ability to modulate (e.g. increase or inhibit) the activity or expression of RYZN. More specifically, the present invention relates to methods of testing compounds for their ability either to increase or to  
25 decrease activity of RYZN. The assays are performed *in vitro* or *in vivo*.

The present invention relates to a method for the screening of a candidate substance for interaction with a polypeptide comprising RYZN extracellular domain, said method comprising the following steps:

- a) providing said polypeptide comprising RYZN extracellular domain;
- 30 b) obtaining a candidate substance;
- c) bringing into contact said polypeptide with said candidate substance;
- d) detecting the complexes formed between said polypeptide and said candidate substance.

The invention further relates to a method for the production of a pharmaceutical composition comprising a method for the screening of a candidate substance that interact with a RYZN polypeptide, fragments or variants thereof and furthermore mixing the identified substance with a pharmaceutically acceptable carrier.

5       The present invention relates to a method for the screening of a candidate substance for the capacity to increase expression of RYZN, said method comprising the following steps:

- a) isolating mRNA from cells which have or have not been contacted with said candidate substance;
- b) carrying out a Northern blot analysis with labeled cDNA probe encoding all or part  
10 of RYZN polypeptide;
- c) wherein increased signal in cells having been contacted with said candidate substance over that of uncontacted cells is taken to indicate that said candidate substance increases expression of RYZN and is an AGONIST of RYZN activity; and
- 15 d) wherein decreased signal in cells having been contacted with said candidate substance over that of uncontacted cells is taken to indicate that said candidate substance decreases expression of RYZN and is an ANTAGONIST of RYZN activity.

Methods of isolating mRNA and carrying out Northern blot analysis are well known to those of  
20 ordinary skill in the art.

#### Preparation of Antibody Compositions

Substantially pure protein or polypeptide is isolated from transfected or transformed cells containing an expression vector encoding the RYZN protein or a portion thereof. The concentration of protein in the final preparation is adjusted, for example, by concentration on an Amicon filter  
25 device, to the level of a few micrograms/ml. Monoclonal or polyclonal antibody to the protein can then be prepared by methods well known to those of ordinary skill in the art.

Preferably the present invention includes monoclonal and polyclonal antibodies that specifically bind RYZN polypeptide fragment comprising the extracellular domain of mature RYZN polypeptide. Particularly preferred soluble fragment of RYZN comprises amino acids 1-61, 2-61, 1-  
30 63, 2-63, 1-68, 2-68, 1-73, 2-73, 1-74, 2-74, 1-76, 2-76, 1-89, 2-89, 1-96, 2-96, 1-108, 2-108, 1-113 or 2-113 of SEQ ID NO:2.

#### EXAMPLES

The following Examples are provided for illustrative purposes and not as a means of limitation. One of ordinary skill in the art would be able to design equivalent assays and methods based on the disclosure herein all of which form part of the instant invention.

EXAMPLE 1: Use of Biacore Technology to Detect Specific Binding of a Test Compound  
5 to Polypeptide Fragment Comprising RYZN Extracellular Domain

Biacore utilizes a biosensor technology for monitoring interactions between two or more molecules in real time, without the use of labels. The molecular classes that can be studied are diverse, ranging from proteins, peptides, nucleic acids, carbohydrates, and lipids to low molecular weight substances and pharmaceuticals.

10 The detection principle is based on the optical phenomena of surface plasmon resonance, which detects changes in refractive index close to a biosensor surface. In a typical experiment one of the interacting molecules is immobilized or captured (here, polypeptide fragment comprising RYZN extracellular domain) to a flexible dextran layer close to the sensor surface. The interacting partner (here, test compound) is flowed across that surface. If an interaction occurs between the two  
15 molecules, there is a resulting increase in signal due to the increase in mass at the chip surface.

Soluble polypeptide fragment comprising RYZN extracellular domain is attached to the sensor surface via amine coupling chemistry. The dextran is activated using N-hydroxysuccinimide and N-ethyl-N'-(3-dimethylaminopropyl)-carbodiimide hydrochloride for 7 minutes. Said RYZN polypeptide fragment is diluted in 10mM Na Acetate pH 5.0 at a concentration of 10µg/ml and  
20 injected over the activated surface for 7 minutes. The surface is then blocked for 7 minutes using ethanolamine to remove any remaining esters. A blank flow cell absent said RYZN polypeptide fragment is set up in parallel and used as a control surface. The running buffer is HBS-EP (0.01M HEPES pH 7.4, 0.15M NaCl, 3mM EDTA, 0.005% Surfactant P20) and the instrument temperature is 25°C.

25 The test compound is filtered through an Ultrafree-0.5 Centrifugal Filter Device and resuspended in HBS-EP running buffer. The test compound is then diluted 1:10 in HBS-EP and injected over the said RYZN polypeptide fragment surface and the blank control surface for 1 minute at a flow rate of 50 µl/min. The sensorgrams from the receptor surface and the control surface are aligned and overlaid.

30 To obtain the specific binding, the control surface was subtracted from the active surface comprised of said RYZN polypeptide fragment.

Example 2: Effect of LIGAND on Muscle Cell Fatty Acid Oxidation

C2C12 cells are differentiated in the presence or absence of 2 µg/mL LIGAND for 4 days. On day 4, oleate oxidation rates are determined by measuring conversion of 1-<sup>14</sup>C-oleate (0.2 mM)



to  $^{14}\text{CO}_2$  for 90 min. This experiment can be used to screen for active polypeptides and peptides as well as AGONISTS and ANTAGONISTS or activators and inhibitors of LIGAND receptor.

The effect of LIGAND on the rate of oleate oxidation can be compared in differentiated C2C12 cells (murine skeletal muscle cells; ATCC, Manassas, VA CRL-1772) and in a hepatocyte cell line (Hepa1-6; ATCC, Manassas, VA CRL-1830). Cultured cells are maintained according to manufacturer's instructions. The oleate oxidation assay is performed as previously described (Muoio et al (1999) Biochem J 338;783-791). Briefly, nearly confluent myocytes are kept in low serum differentiation media (DMEM, 2.5% Horse serum) for 4 days, at which time formation of myotubes became maximal. Hepatocytes are kept in the same DMEM medium supplemented with 10% FCS for 2 days. One hour prior to the experiment the media is removed and 1 mL of preincubation media (MEM, 2.5% Horse serum, 3 mM glucose, 4 mM Glutamine, 25 mM Hepes, 1% FFA free BSA, 0.25 mM Oleate, 5  $\mu\text{g/mL}$  gentamycin) is added. At the start of the oxidation experiment  $^{14}\text{C}$ -Oleic acid (1 $\mu\text{Ci/mL}$ , American Radiolabelled Chemical Inc., St. Louis, MO) is added and cells are incubated for 90 min at 37°C in the absence/presence of 2.5  $\mu\text{g/mL}$  LIGAND. After the incubation period 0.75 mL of the media is removed and assayed for  $^{14}\text{C}$ -oxidation products as described below for the muscle FFA oxidation experiment.

#### EXAMPLE 3: Effect of LIGAND on *In Vitro* Glucose Uptake by Muscle Cells

L6 Muscle cells are obtained from the European Culture Collection (Porton Down) and are used at passages 7-11. Cells are maintained in standard tissue culture medium DMEM, and glucose uptake is assessed using [ $^3\text{H}$ ]-2-deoxyglucose (2DG) with or without LIGAND in the presence or absence of insulin ( $10^{-8}$  M) as has been previously described (Walker, P.S. et al. (1990) Glucose transport activity in L6 muscle cells is regulated by the coordinate control of subcellular glucose transporter distribution, biosynthesis, and mRNA transcription. JBC 265(3):1516-1523; and Kilp, A. et al. (1992) Stimulation of hexose transport by metformin in L6 muscle cells in culture. Endocrinology 130(5):2535-2544, which disclosures are hereby incorporated by reference in their entireties). Uptake of 2DG is expressed as the percentage change compared with control (no added insulin or LIGAND). Values are presented as mean  $\pm$  SEM of sets of 4 wells per experiment. Differences between sets of wells are evaluated by Student's t test, probability values  $p < 0.05$  are considered to be significant.

#### EXAMPLE 4: Effect of LIGAND on Mice Fed a High-Fat Diet

Experiments are performed using approximately 6 week old C57Bl/6 mice (8 per group). All mice are housed individually. The mice are maintained on a high fat diet throughout each experiment. The high fat diet (cafeteria diet; D12331 from Research Diets, Inc.) has the following composition: protein kcal% 16, sucrose kcal% 26, and fat kcal% 58. The fat is primarily composed of coconut oil, hydrogenated.

After the mice are fed a high fat diet for 6 days, micro-osmotic pumps are inserted using isoflurane anesthesia, and are used to provide LIGAND, saline, and an irrelevant peptide to the mice

subcutaneously (s.c.) for 18 days. LIGAND is provided at doses of 100, 50, 25, and 2.5 µg/day and the irrelevant peptide is provided at 10 µg/day. Body weight is measured on the first, third and fifth day of the high fat diet, and then daily after the start of treatment. Final blood samples are taken by cardiac puncture and are used to determine triglyceride (TG), total cholesterol (TC), glucose, leptin, and insulin levels. The amount of food consumed per day is also determined for each group.

EXAMPLE 5: Effect of LIGAND on Plasma Free Fatty Acid in C57 BL/6 Mice

The effect of LIGAND on postprandial lipemia (PPL) in normal C57BL6/J mice is tested.

The mice used in this experiment are fasted for 2 hours prior to the experiment after which a baseline blood sample is taken. All blood samples are taken from the tail using EDTA coated capillary tubes (50 µL each time point). At time 0 (8:30 AM), a standard high fat meal (6g butter, 6 g sunflower oil, 10 g nonfat dry milk, 10 g sucrose, 12 mL distilled water prepared fresh following Nb#6, JF, pg.1) is given by gavage (vol.=1% of body weight) to all animals.

Immediately following the high fat meal, 25µg a LIGAND is injected i.p. in 100 µL saline. The same dose (25µg/mL in 100µL) is again injected at 45 min and at 1 hr 45 min. Control animals are injected with saline (3x100µL). Untreated and treated animals are handled in an alternating mode.

Blood samples are taken in hourly intervals, and are immediately put on ice. Plasma is prepared by centrifugation following each time point. Plasma is kept at -20°C and free fatty acids (FFA), triglycerides (TG) and glucose are determined within 24 hours using standard test kits (Sigma and Wako). Due to the limited amount of plasma available, glucose is determined in duplicate using pooled samples. For each time point, equal volumes of plasma from all 8 animals per treatment group are pooled.

EXAMPLE 6: Effect of LIGAND on Plasma FFA, TG and Glucose in C57 BL/6 Mice

Briefly, 14 mice are fasted for 2 hours prior to the experiment after which a baseline blood sample is taken. All blood samples are taken from the tail using EDTA coated capillary tubes (50 µL each time point). At time 0 (9:00AM), a standard high fat meal (see Example 6) is given by gavage (vol.=1% of body weight) to all animals. Immediately following the high fat meal, 4 mice are injected 25 µg of LIGAND i.p. in 100µL saline. The same dose (25µg in 100µL) is again injected at 45 min and at 1 hr 45 min. A second treatment group receives 3 times 50 µg LIGAND at the same intervals. Control animals are injected with saline (3x100µL). Untreated and treated animals are handled in an alternating mode.

Blood samples are immediately put on ice. Plasma is prepared by centrifugation following each time point. Plasma is kept at -20 °C and free fatty acids (FFA), triglycerides (TG) and glucose are determined within 24 hours using standard test kits (Sigma and Wako).

EXAMPLE 7: Effect of LIGAND on FFA following Epinephrine Injection

In mice, plasma free fatty acids increase after intragastric administration of a high fat/sucrose test meal. These free fatty acids are mostly produced by the activity of lipolytic enzymes

*i.e.* lipoprotein lipase (LPL) and hepatic lipase (HL). In this species, these enzymes are found in significant amounts both bound to endothelium and freely circulating in plasma. Another source of plasma free fatty acids is hormone sensitive lipase (HSL) that releases free fatty acids from adipose tissue after  $\beta$ -adrenergic stimulation. To test whether LIGAND also regulates the metabolism of free fatty acid released by HSL, mice are injected with epinephrine.

Two groups of mice are given epinephrine (5 $\mu$ g) by intraperitoneal injection. A treated group is injected with a LIGAND (25 $\mu$ g) one hour before and again together with epinephrine, while control animals receive saline. Plasma is isolated and free fatty acids and glucose are measured.

EXAMPLE 8: Effect of LIGAND on FFA following Intralipid Injection

Two groups of mice are intravenously (tail vein) injected with 30  $\mu$ L bolus of Intralipid-20% (Clintec) to generate a sudden rise in plasma FFAs, thus by-passing intestinal absorption. (Intralipid is an intravenous fat emulsion used in nutritional therapy). A treated group (LIGAND-treated) is injected with LIGAND (25 $\mu$ g) at 30 and 60 minutes before Intralipid is given, while control animals receive saline. Plasma is isolated and FFAs are measured as described previously. The effect of LIGAND on the decay in plasma FFAs following the peak induced by Intralipid injection is then monitored.

EXAMPLE 9: Effect of LIGAND on Weight Gain and Weight Loss of Mice and on Maintenance of Weight Loss in Mice

In the first experiment, 10-week-old male C57BL/6J mice are put on a very high fat/sucrose purified diet for 19 days to promote weight gain; the average body weight at this time is 30g. The mice are then surgically implanted with an osmotic pump (Alzet, Newark, DE) delivering either 2.5  $\mu$ g/day of LIGAND or physiological saline. The mice are continued on the high fat diet and their body weight was recorded over the following 10-day period.

Weight gain by mice treated with saline in contradistinction to weight loss by mice treated with LIGAND is taken as evidence that in this inbred strain of normal mice, a continuous infusion of a daily low dose of LIGAND can prevent weight gain caused by high fat/sucrose feeding, in a sustainable way.

Data are expressed throughout as mean  $\pm$  SEM; a p-value  $< 0.05$  is considered statistically significant. Statistical analysis is typically done using either the unpaired Student's t test or the paired Student's t test.

Maintenance of weight loss in mice

In order to demonstrate the ability of LIGAND to maintain weight loss, normal mice are put on a reduced calorie diet to promote weight loss. The reduced calorie diet is continued until the mice lose 10% of their initial weight. A second group of mice are continued on the reduced calorie diet until the mice lose 20% of their initial weight. The mice are then surgically implanted with an osmotic pump (Alzet, Newark, DE) delivering either 2.5  $\mu$ g/day of LIGAND or physiological saline. The mice are returned to a normal diet and their body weights are recorded over a 10-day period.

After 10 days, the outcome wherein mice treated with LIGAND have a lower weight than mice treated with saline is taken to provide evidence that treatment with LIGAND promotes the maintenance of weight loss.

EXAMPLE 10: Assessment of homotrimer formation by gAPM1, gC2P, gZADJ-2 or gZADJ-7

5 polypeptide fragment.

Homotrimer formation by gAPM1, gC2P, gZADJ-2 or gZADJ-7 polypeptide fragment is assessed using sedimentation equilibrium in analytical centrifuges, a method that determines molecular weight accurately and independently of other physical factors such as shape.

- Candidate gAPM1, gC2P, gZADJ-2 or gZADJ-7 polypeptide fragment homotrimer is
- 10 purified, for example using a protocol comprising a method of gel filtration such as 16/60 superdex 200 gel filtration column (Amersham). Said purified candidate gAPM1, gC2P, gZADJ-2 or gZADJ-7 polypeptide fragment homotrimer protein concentration is made 3  $\mu$ M in 5.7 mM phosphate (pH 7.5), 137 mM NaCl, 2.7 mM KCl. Samples are centrifuged at 8,000 rpm for 18 hours at 10°C in a Beckman XL-A analytical ultracentrifuge before absorbance is recorded. The data are fit globally,
- 15 using MacNonlin PPC [Johnson ML et al., Biophys J (1981) 36:575-8; Schuster TM et al., Curr Opin Struct Biol (1996) 6:650-8; Hensley P, Structure (1996) 4:367-73; the disclosures of which are incorporated herein by reference in their entirety] to the following equation that describes the sedimentation of a homogeneous species:  $Abs = B + A' \exp[H \times M (x^2 - x_0^2)]$  where Abs = absorbance at radius x, A' = absorbance at reference radius  $x_0$ ,  $H = (1 - \nu\rho)\omega^2/2RT$ , R = gas constant, T =
- 20 temperature in Kelvin,  $\nu$  = partial specific volume = 0.71896131 mL/g,  $\rho$  = density of solvent = 1.0061 g/mL,  $\omega$  = angular velocity in radians/s, M = apparent molecular weight, and B = solvent absorbance (blank).

TABLE 1

Amino Acid Residues Comprising the Structural Domains of RYZN

SEQ ID NO: 2 Description

EC DOMAIN	TRANSMEMBRANE DOMAIN	IC DOMAIN
1-113	114-136	137-247

.5

EC, extracellular domain; IC, intracellular domain

10

## APM1, C2P, ZADJ-2 and ZADJ-7

>APM1 polypeptide sequence:

5 MLLLGAVLLLLALPGHDQETTTQGPVLLPLPKGACTGWMAGIPGHPGHNGAPGRDGRD  
GTPGEKGEKGDPLIGPKGDIGETGVPGAEGPRGFPGIQGRKGEPGEGAYVYRSAFSVGLET  
YVTIPNMPIRFTKIFYNQONHYDGSTGKFHCNIPGLYYFAYHITVYMKDVKVSLFKKDKAM  
LFTYDOYOENNVDQASGSVLLHLEVGDQVWLOVYGEGERNGLYADNDNDSTFTGFLLYH  
DTN (244)

10

> C2P polypeptide sequence:

MRIWWLLLAIEICTGNINSQDTCRQHPGIPGNPGHNGLPGRDGRDGA KGDKGDAGEPG  
RPGSPGKDGTSGEKGGERGADGKVEAKGIKGDQGSRGSPGKHGPKGLAGPMGEKGLRGETG  
PQGQKGNKGDVGPTGPEGPRGNIGPLGPTGLPGPMGPIGKPGPKGEAGPTGPQGE PGVRGIR  
15 GWKGD RGEKGKIGETLVLPKSAFTVGLTVLSKFPSSDMPIKFDKILYNEFNHYDTAAGKFTC  
HIAGVYYFTYHITVFSRNVQVSLVKNGVKILHTKDAYMSSDQASGGIVLOLKLGDDEVWLO  
VTGGERFNGLEADEDDDTTFTGFLLFSSP (333)

>ZADJ-2 polypeptide sequence:

20 MIPWVLLACALPCAADPLLGA FARRDFRKGSPQLVCSLPGPQGPPGPPGAPGPSGMMGRM  
GFPGKDGQDGHGDRGDSGEEGPPGRTGNRGKPGPKGKAGAI GRAGPRGPKGVNGTPGK  
HGTPGKKGPKGKKGEPLPGPCSCSGHTKSAFSVA VTKSYPRERLPIKFDKILMNEGGHY  
NASSGKFVCGVPGIYYFTYDITLANKHLAIGLVHNGOYRIRTFDANTGNH DVASGSTILALK  
QGDEVWLOIFYSEONGLEYDPYWTDSLFTGFLIYADQDDPNEV (285)

25

>ZADJ-7 polypeptide sequence:

MGKEDTQETRTEPKMFVLLYVTSFAICASGQPRGNQLKGENYSPRYICSIPGLPGPPGPPG  
ANGSPGPHGRIGLPGRDGRDGRKG EKGEKGTAGLRGKTGPLGLAGEKGDQGETGKKGPIG  
PEGEKGEVGPIGPPGPKGDRGEQGDPLPGVCRCGSIVLKSASFVGITTSYPEERLP IIFNKVL  
30 FNEGEHYNPATGKFICAFPGIYYFSYDITLANKHLAIGLVHNGOYRIKTFDANTGNH DVASG  
STVIYLOPEDEVWLEIFFTDQNGLESDPGWADSLFSGFLLYVDTDYLD SISEDDEL (303)

What is claimed is:

1. A method of screening for an AGONIST or an ANTAGONIST of RYZN activity.
2. The method of Claim 1, wherein said activity is selected from the group consisting  
5 of lipid partitioning, lipid metabolism, insulin-like activity, free fatty acid oxidation, and weight reduction.
3. An AGONIST or an ANTAGONIST of RYZN activity.
4. The AGONIST or the ANTAGONIST of Claim 3, wherein said activity is selected  
from the group consisting of lipid partitioning, lipid metabolism, insulin-like activity, free fatty acid  
10 oxidation, and weight reduction.
5. A pharmaceutical or physiologically acceptable composition comprising, consisting  
essentially of, or consisting of the AGONIST or the ANTAGONIST of Claim 3.
6. A method of preventing or treating an obesity-related disease or disorder comprising  
providing or administering to an individual in need of such treatment the composition of Claim 5.

## SEQUENCE LISTING

&lt;110&gt; Genset

&lt;120&gt; TREATMENT OF METABOLIC DISORDERS WITH RYZN AGONISTS AND ANTAGONISTS

&lt;130&gt; 163.WO1

&lt;150&gt; 60/334,152

&lt;151&gt; 2001-11-28

&lt;160&gt; 2

&lt;170&gt; FastSEQ for Windows Version 4.0

&lt;210&gt; 1

&lt;211&gt; 1192

&lt;212&gt; DNA

&lt;213&gt; Homo sapiens

&lt;220&gt;

&lt;221&gt; CDS

&lt;222&gt; (6)...(749)

&lt;400&gt; 1

gagta atg agt ggc ctg ggc cgg agc agg cga ggt ggc cgg agc cgt gtg 50

Met Ser Gly Leu Gly Arg Ser Arg Arg Gly Gly Arg Ser Arg Val

1 5 10 15

gac cag gag gag cgc tgg tca ctc agc tgc cgc aag gag caa ggc aag 98

Asp Gln Glu Glu Arg Trp Ser Leu Ser Cys Arg Lys Glu Gln Gly Lys

20 25 30

ttc tat gac cat ctc ctg agg gac tgc atc agc tgt gcc tcc atc tgt 146

Phe Tyr Asp His Leu Leu Arg Asp Cys Ile Ser Cys Ala Ser Ile Cys

35 40 45

gga cag cac cct aag caa tgt gca tac ttc tgt gag aac aag ctc agg 194

Gly Gln His Pro Lys Gln Cys Ala Tyr Phe Cys Glu Asn Lys Leu Arg

50 55 60

agc cca gtg aac ctt cca cca gag ctc agg aga cag cgg agt gga gaa 242

Ser Pro Val Asn Leu Pro Pro Glu Leu Arg Arg Gln Arg Ser Gly Glu

65 70 75

gtt gaa aac aat tca gac aac tcg gga agg tac caa gga ttg gag cac 290

Val Glu Asn Asn Ser Asp Asn Ser Gly Arg Tyr Gln Gly Leu Glu His

80 85 90 95

aga ggc tca gaa gca agt cca gct ctc ccg ggg ctg aag ctg agt gca 338

Arg Gly Ser Glu Ala Ser Pro Ala Leu Pro Gly Leu Lys Leu Ser Ala

100 105 110

gat cag gtg gcc ctg gtc tac agc acg ctg ggg ctc tgc ctg tgt gcc 386

Asp Gln Val Ala Leu Val Tyr Ser Thr Leu Gly Leu Cys Leu Cys Ala

115 120 125

gtc ctc tgc tgc ttc ctg gtg gcg gtg gcc tgc ttc ctc aag aag agg 434



2

Val Leu Cys Cys Phe Leu Val Ala Val Ala Cys Phe Leu Lys Lys Arg  
 130 135 140

ggg gat ccc tgc tcc tgc cag ccc cgc tca agg ccc cgt caa agt ccg 482  
 Gly Asp Pro Cys Ser Cys Gln Pro Arg Ser Arg Pro Arg Gln Ser Pro  
 145 150 155

gcc aag tct tcc cag gat cac gcg atg gaa gcc ggc agc cct gtg agc 530  
 Ala Lys Ser Ser Gln Asp His Ala Met Glu Ala Gly Ser Pro Val Ser  
 160 165 170 175

aca tcc ccc gag cca gtg gag acc tgc agc ttc tgc ttc cct gag tgc 578  
 Thr Ser Pro Glu Val Glu Thr Cys Ser Phe Cys Phe Pro Glu Cys  
 180 185 190

agg gcg ccc acg cag gag agc gca gtc acg cct ggg acc ccc gac ccc 626  
 Arg Ala Pro Thr Gln Glu Ser Ala Val Thr Pro Gly Thr Pro Asp Pro  
 195 200 205

act tgt gct gga agg tgg ggg tgc cac acc agg acc aca gtc ctg cag 674  
 Thr Cys Ala Gly Arg Trp Gly Cys His Thr Arg Thr Thr Val Leu Gln  
 210 215 220

cct tgc cca cac atc cca gac agt ggc ctt ggc att gtg tgt gtg cct 722  
 Pro Cys Pro His Ile Pro Asp Ser Gly Leu Gly Ile Val Cys Val Pro  
 225 230 235

gcc cag gag ggg ggc cca ggt gca taa atgggggtca gggaggga 769  
 Ala Gln Glu Gly Gly Pro Gly Ala \*  
 240 245

ggaggagggga gagagatgga gaggagggga gagagaaaga gaggtgggga gaggggagag 829  
 agatatgagg agagagagac agaggaggca gagagggaga gaaacagagg agacagagag 889  
 ggagagagag acagaggggag agagagacag aggggaagag aggcagagag ggaaagaggc 949  
 agagaaggaa agaggcagag agggagagag gcagagaggg agagaggcag agagacagag 1009  
 agggagagag ggacagagag agatagagca ggaggtcggg gcactctgag tcccagttcc 1069  
 cagtgcagct gtaggtcgtc atcacctaac cacacgtgca ataaagtcct cgtgcctgct 1129  
 gctcacagcc cccgagagcc cctcctcctg gagaataaaa cctttggcag ctgcccttcc 1189  
 tca 1192

<210> 2  
 <211> 247  
 <212> PRT  
 <213> Homo sapiens

<400> 2  
 Met Ser Gly Leu Gly Arg Ser Arg Arg Gly Gly Arg Ser Arg Val Asp  
 1 5 10 15  
 Gln Glu Glu Arg Trp Ser Leu Ser Cys Arg Lys Glu Gln Gly Lys Phe  
 20 25 30  
 Tyr Asp His Leu Leu Arg Asp Cys Ile Ser Cys Ala Ser Ile Cys Gly  
 35 40 45  
 Gln His Pro Lys Gln Cys Ala Tyr Phe Cys Glu Asn Lys Leu Arg Ser  
 50 55 60  
 Pro Val Asn Leu Pro Pro Glu Leu Arg Arg Gln Arg Ser Gly Glu Val  
 65 70 75 80  
 Glu Asn Asn Ser Asp Asn Ser Gly Arg Tyr Gln Gly Leu Glu His Arg  
 85 90 95

3

Gly	Ser	Glu	Ala	Ser	Pro	Ala	Leu	Pro	Gly	Leu	Lys	Leu	Ser	Ala	Asp
		100						105						110	
Gln	Val	Ala	Leu	Val	Tyr	Ser	Thr	Leu	Gly	Leu	Cys	Leu	Cys	Ala	Val
		115						120						125	
Leu	Cys	Cys	Phe	Leu	Val	Ala	Val	Ala	Cys	Phe	Leu	Lys	Lys	Arg	Gly
		130						135						140	
Asp	Pro	Cys	Ser	Cys	Gln	Pro	Arg	Ser	Arg	Pro	Arg	Gln	Ser	Pro	Ala
145						150						155		160	
Lys	Ser	Ser	Gln	Asp	His	Ala	Met	Glu	Ala	Gly	Ser	Pro	Val	Ser	Thr
				165						170				175	
Ser	Pro	Glu	Pro	Val	Glu	Thr	Cys	Ser	Phe	Cys	Phe	Pro	Glu	Cys	Arg
		180						185						190	
Ala	Pro	Thr	Gln	Glu	Ser	Ala	Val	Thr	Pro	Gly	Thr	Pro	Asp	Pro	Thr
		195						200						205	
Cys	Ala	Gly	Arg	Trp	Gly	Cys	His	Thr	Arg	Thr	Thr	Val	Leu	Gln	Pro
210						215						220			
Cys	Pro	His	Ile	Pro	Asp	Ser	Gly	Leu	Gly	Ile	Val	Cys	Val	Pro	Ala
225						230						235		240	
Gln	Glu	Gly	Gly	Pro	Gly	Ala									
				245											

## INTERNATIONAL SEARCH REPORT

Internati

location No

PCT/IB 02/04581

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 A61K38/17 A61K45/00 G01N33/50 A61P3/00

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 A61K A61P G01N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ, BIOSIS, MEDLINE, SEQUENCE SEARCH

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 01 51645 A (GENSET SA ;YEN POTIN FRANCES (US); BIHAIN BERNARD (US); FRUEBIS JO) 19 July 2001 (2001-07-19) claims page 58, line 11 -page 59, line 15 ---	1-6
X	EP 1 033 134 A (OTSUKA PHARMA CO LTD) 6 September 2000 (2000-09-06) page 2, line 44 - line 47 ---	1-6
X	WO 00 68380 A (INCYTE GENOMICS INC ;AZIMZAI YALDA (US); YUE HENRY (US); BANDMAN O) 16 November 2000 (2000-11-16) SEQ ID NO 8 page 35, line 17 -page 37, line 5 --- -/--	1-6



Further documents are listed in the continuation of box C.



Patent family members are listed in annex.

## \* Special categories of cited documents :

\*A\* document defining the general state of the art which is not considered to be of particular relevance

\*E\* earlier document but published on or after the international filing date

\*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

\*O\* document referring to an oral disclosure, use, exhibition or other means

\*P\* document published prior to the international filing date but later than the priority date claimed

\*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

\*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

\*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

\*&amp;\* document member of the same patent family

Date of the actual completion of the international search

17 January 2003

Date of mailing of the international search report

27/01/2003

Name and mailing address of the ISA

European Patent Office, P.B. 5618 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Böhmerova, E

## INTERNATIONAL SEARCH REPORT

Internatio plication No

PCT/IB 02/04581

## C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	WO 99 59618 A (SMITHKLINE BEECHAM CORP) 25 November 1999 (1999-11-25) page 1, line 23 - line 32 page 15, line 17 - line 20 -----	1-6
X	WO 00 73448 A (ZYMOGENETICS INC) 7 December 2000 (2000-12-07) page 86, line 21 -page 88, line 23 page 72, line 28 -page 73, line 33 -----	1-6
X	WO 00 40716 A (ZYMOGENETICS INC) 13 July 2000 (2000-07-13) SEQ ID NOs 2, 4 page 47, line 13 -page 52, line 27 page 57, line 3 - line 7 claims 29-63 -----	1-6

## INTERNATIONAL SEARCH REPORT

International Publication No  
PCT/IB 02/04581

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
WO 0151645	A	19-07-2001	AU 5789901 A BR 0107626 A EP 1248849 A1 WO 0151645 A1 US 2002058617 A1 US 2002037849 A1 US 2002091080 A1	24-07-2001 15-10-2002 16-10-2002 19-07-2001 16-05-2002 28-03-2002 11-07-2002
EP 1033134	A	06-09-2000	EP 1033134 A1 US 6461821 B1 WO 9921577 A1	06-09-2000 08-10-2002 06-05-1999
WO 0068380	A	16-11-2000	AU 5129800 A EP 1177296 A2 WO 0068380 A2	21-11-2000 06-02-2002 16-11-2000
WO 9959618	A	25-11-1999	EP 1085897 A1 EP 1083923 A1 JP 2002515452 T WO 9959618 A1 WO 9959619 A1 US 2001009905 A1	28-03-2001 21-03-2001 28-05-2002 25-11-1999 25-11-1999 26-07-2001
WO 0073448	A	07-12-2000	AU 5159500 A EP 1185645 A1 WO 0073448 A1	18-12-2000 13-03-2002 07-12-2000
WO 0040716	A	13-07-2000	AU 2408400 A BG 105674 A BR 0007423 A CN 1342202 T CZ 20012465 A3 EE 200100360 A EP 1141274 A2 HU 0200429 A2 NO 20013316 A SK 9712001 A3 TR 200101998 T2 WO 0040716 A2	24-07-2000 29-03-2002 22-01-2002 27-03-2002 13-03-2002 16-12-2002 10-10-2001 29-06-2002 06-09-2001 05-02-2002 21-06-2002 13-07-2000